

## **Influence of Resistance Training on Anabolic Hormones in Pre-Pubertal and Pubertal Males**

**Tsolakis, Charilaos\* and Bogdanis, Gregory C. \*\***

\*Department of Track & Field and \*\*Department of Sports Medicine & Biology of Physical Activity, Faculty of Physical Education and Sports Science, University of Athens, Greece

Corresponding Author: Tsolakis Charilaos, Ph.D, 6 Kotsika str. 10434 Athens, Greece, Email: tsolakis@phed.uoa.gr

### **Abstract**

This review focuses on the hormonal responses to resistance training in pre-pubertal and pubertal males with the aim to elucidate possible mechanisms that may be responsible for the increases in strength and power following systematic training in these age groups. Studies that have controlled for the confounding effects of growth and motor skill acquisition, have shown that resistance training results in significant increases in muscular strength even before puberty. Although the training induced muscle hypertrophy in pre-pubescent is possible, it is much less compared that of post-pubescent. On the other hand an equal or greater improvement in neuromuscular activation, motor skill and intrinsic muscular adaptations are reported in the pre-pubescent compared to the post-pubescent following resistance training programs. Resting serum testosterone concentration has been found to increase following 2-12 month resistance training programs and this is accompanied with an increase in muscle strength. Changes in strength and hormonal concentrations depend on the characteristics of the resistance training program, with the most important being exercise intensity and volume.

**Key Words: Growth Hormone, Testosterone, Children**

### **Introduction**

Resistance training for children and adolescents has been a topic of great interest among scientists, physicians, coaches, young athletes and their parents. The research conducted in this area during the last two decades has provided valuable information on the responses of the young organism to this type of training and has advanced our understanding on mechanisms responsible for these adaptations (*Falk and Tenenbaum, 1996*). This review focuses on the hormonal responses to resistance training in pre-pubertal and pubertal males with the aim to elucidate possible mechanisms that may be responsible for the increases in strength and power following systematic training in these age groups.

### **Effectiveness of resistance training in pre-pubertal and pubertal males**

The participation of boys in resistance training programs leads to a positive

influence on their fitness that results in an increase in sport performance and offers protection from sports injuries (*Kraemer-Fleck, 1993*). Resistance training in young boys also improves quality of life and contributes to positive attitude toward wellness and exercise (*Shephard, 1984*). One of the most important characteristics of resistance exercise programs for children must be safety. Emphasis should be placed on proper technique and sessions should be supervised by a qualified instructor. Exercise intensity and interval between exercises should be chosen according to the maturation level of the young athlete, but maximal loading should be avoided (*Bases, 2004*).

Heavy resistance exercise in adults' results in significant adaptive responses, including increases in strength and power and muscle hypertrophy that, are dependent on the type, intensity of loading and volume of training (*Fleck and Kraemer, 1987*). It is well known that in adults, the increases in

strength during the early phase of training are attributed mainly to neural factors, but as training proceeds, there is a gradually increased contribution of muscular hypertrophy to the increases in strength (Moritani, 1992). The effectiveness of resistance training in children seems to be dependent primarily on the provision of a sufficient training intensity and volume and to a lesser degree on training duration, following the principle of specificity. Strength increases ranging from 5 to 50% have been reported using isometric, isotonic and isokinetic training methods involving high intensity loads (Kraemer et al., 1989, Sale, 1989). As training progresses there is an interaction between neural factors and muscle hypertrophy. In pre-pubescent children, neural factors would predominate, but as the child reaches puberty the growth and maturation related hormones favor muscle hypertrophy (Kraemer et al., 1989, Sale, 1989). The optimal combination of training method, mode, intensity, volume and duration of training for maximal strength gain during preadolescence however, has yet to be determined.

Irrespective of the mechanism responsible, muscular strength gains have been evident across all maturity stages, including pre-pubescents (Nielsen et al., 1980, Weltman et al., 1986, Blimkie et al., 1989, Ramsay et al., 1990, Hassan, 1991, Fukunaga et al., 1992, Faigenbaum et al., 1993, Ozmun et al., 1994), pubescents and post pubescent individuals (Pfeifer-Francis, 1986, Sailors-Berg, 1987, Sale, 1989). Studies that have controlled for the confounding effects of growth and motor skill acquisition, have shown that resistance training results in substantial and significant increases in strength before puberty beyond what would occur due to the normal growth (Nielsen et al., 1980, Weltman et al., 1986, Sailors-Berg, 1987,

Hassan, 1991, Fukunaga et al., 1992, Ozmun et al., 1994). The trainability of strength during the post-pubertal period is a less contentious issue. Significant strength gains in this age group have resulted from isometric training, dynamic or isotonic weight training, isokinetic and hydraulic resistance training and appear to be directly related to training frequency (Blimkie & Bar-Or, 1996).

The conclusions from the studies which examine the effectiveness of resistance training during preadolescence compared to training during adolescence and adulthood are still debatable (Nielsen et al., 1980, Pfeifer and Francis, 1986, Sailors and Berg, 1987, Sale, 1989). Studies often suffer from methodological shortcomings, such as absence of a control group, inappropriate training program and lack of statistical power to allow for sound conclusions (Falk and Tenenbaum, 1996). Nevertheless, most, but not all, studies indicate that when the relative training loads are moderate to high and the groups are formulated according to distinct maturity levels, it appears that pre-pubescents are probably less trainable in terms of absolute strength, but equally if more trainable in percentage improvements compared to adolescents and adults (Hakkinen et al., 1989, Sailors and Berg, 1987, Sale, 1989, Fukunaga et al., 1992). Although the training induced muscle hypertrophy in pre-pubescents is possible (Merish and Stoboy, 1989, Fukunaga et al., 1992), it is much less compared that of post-pubescents. On the other hand an equal or greater improvement in neuromuscular activation, motor skill and intrinsic muscular adaptations are reported in the pre-pubescents compared to the post-pubescents following resistance training programs. (Blimkie and Bar-Or, 1996)

### **Hormonal mechanisms influencing gains in muscular strength**

The pubertal growth spurt is influenced by the release of important hormones such as Growth Hormone (GH), insulin-like growth factor I (IGF-I) and steroid sex hormones that induce increases in growth velocity, bone and muscle maturation, functional ability and several metabolic adaptations (Rogol, 1994). These changes may influence the development of physical capacity and performance during childhood and adolescence. Adrenarche (as defined by synthesis and secretion of androgens when the adrenal zona reticularis matures) is characterized by a progressive increase in the secretion of adrenal hormones. The gonadarche (as defined by activation of the testes and ovaries at puberty) follows 2 to 3 years after this first stage and is under the control of GH, IGF-I and steroid sex hormones (Lee, 1995). The release of these hormones is controlled by the pulsatile stimulation of a hypothalamic gonadotrophin – releasing factor. The processes that initiate the onset and progression of human puberty are not completely understood. The dual role of the recently discovered hormone leptin, i.e. as a regulator of fat mass and as a metabolic signal to the reproductive axis, suggests that leptin may be a permissive factor responsible for the onset of puberty (Matzoros et al., 1997).

Increased plasma levels of GH, testosterone, estradiol and progesterone have an anabolic effect on structural protein production. The synergistic action of GH and gonadal steroids promotes the pubertal growth spurt mainly in bones and muscles and this may contribute to metabolic and hormonal responses of children and adolescents during exercise. (Boisseau and Delamarche, 2000). Such mechanisms appear to be operational in response to a

training stimulus provided by heavy resistance exercise and could theoretically influence the growth and remodeling process of various body tissues over the course of a resistance training program.

The endocrine system of the body adapts to the repeated stimulation of exercise training by: a) altering the intensity of the exercise stimulus necessary to increase or decrease hormone secretion, b) altering the tissue responsiveness to the hormone. This is done by increasing or decreasing the number of circulating proteins that bind the hormone and protect it from degradation but also render it biologically inactive, as well as changing the number of cellular hormone receptors or post receptor function, and c) working through neuroendocrine adaptations to alter basal and maximal hormone secretion (Roemmich, 2005). The endocrine environment can have a profound impact on the adaptation process of skeletal muscle to resistance exercise and it is well known that acute as well as long-term resistance training activates a wide variety of neuroendocrine mechanisms (Kraemer, 1988, Hakkinen, 1989, Kraemer, 1992a).


Hormonal variables have also been used as indicators of the magnitude of training stress during resistance exercise and other forms of training (Adlecreutz et al., 1986, Hakkinen et al., 1987, Fry et al., 1993). It has been noted that chronic hormonal adaptations have been correlated to changes in muscular strength and power for competitive Olympic style weight lifters (Hakkinen et al., 1987, Fry et al., 2000), as well as to fiber type alterations in previously untrained men (Staron et al., 1994) and consequently, the various hormonal mechanisms probably respond differentially in trained and untrained individuals (Hakkinen, 1989).

**Role of Testosterone**


Testosterone (T), the male sex hormone, is considered to be an anabolic hormone and contributes also to metabolic control (Viru and Viru, 2005). The “free: or biologically available T hypothesis, supports the idea that only the free hormone is transported to the target tissues, suggesting that the level of free androgen index (FAI) may be of importance for trainability (Kraemer et al 1990). FAI is the ratio of the concentration of T and the sex hormone binding globulin (SHBG), and thus the amount of active T is determined by SHBG concentration (Remes et al, 1979). SHBG may also act as a hormone itself and may have a distinct biological activity, as also suggested for other binding proteins (Rosner, 1991). The SHBG declines progressively during pubertal phases in healthy boys (Anderson, 1974).

Exercise can acutely increase or decrease circulating T, depending on the mode and the intensity of exercise (Schmid et al., 1982). Increases occur during and after relatively short, high intensity work such as resistance training or sprint events, while declines are associated with increasing duration endurance events especially marathon and ultra running events (Kuoppasalmi et al., 1980, Schurmeyer et al., 1984). In resistance training, the main role of T is the induction of synthesis of contractile proteins in involved muscles. Beside that, during acute resistance exercises, as well as during competition, T action seems to be essential for mobilizing performance capacity (Viru and Viru, 2005). Although there is no evidence that T influences the neural adaptations in resistance training, essential is the influence of high preconditioning T levels on central nervous structures. (Kurz et al., 1986, Araki et al., 1991, Matsumoto, 1992)

The initial report of Fahey et al., (1976) supports the possibility that resistance training may alter hypopituitary axis in younger males. However, studies on the responses of T to resistance training of young males have only recently appeared in the literature (Kraemer et al., 1989). The effect of the increase in T in growing individuals due to resistance training on growth and development remains to be studied.



**Figure 1. Resting testosterone (T) and growth hormone (GH) in pre-pubertal male athletes of different sports and in non-athletes (Control). Modified with permission from Tsolakis et al. 2003, Hormones: 2(2): 103-112**



**Figure 2. Sex hormone binding globulin (SHBG) and free androgen index (FAI) in pre-pubertal male athletes of different sports and in non-athletes (Control). Modified with permission from Tsolakis et al. 2003, Hormones: 2(2): 103-112**

### **Role of Growth Hormone**

Growth hormone (GH) has been linked to the promotion of anabolism in both muscle and connective tissue. Specifically, it enhances cellular amino acid uptake and protein synthesis in skeletal muscle, resulting in hypertrophy of both muscle fiber types (Crisis *et al.*, 1991). The anabolic effects of GH result from direct and indirect interactions with androgens (Jorgensen *et al.*, 1996), and thyroid hormone (Weiss and Refetoff, 1996) and are mediated via IGF-I (Florini *et al.*, 1996). The complexity of pituitary production and release of GH and binding proteins is just beginning to be elucidated (Nindl, 2003). It appears that GH responses depend upon type, load and volume of exercise. This suggests that a threshold exist for intensity of exercise to elicit a significant response of the hypopituitary axis to resistance exercise (Kraemer, 1992). Activation of a relatively large muscle mass also appears to be a vital element of the exercise stimulus to elicit a significant GH response. Resistance exercise workouts that have short rest periods (1 min rest between sets and exercises, moderate-to-high intensities (load: 8-12 RM) and whole body workout protocols (8-10 exercises) results in the highest GH changes in the blood (Kraemer *et al.*, 1990, 1991, Hakkinen and Pakarinen, 1995, Roemmich and Rogol 1997, Hansen *et al.*, 2001). Surprisingly, no changes in the resting concentrations of GH have been observed in response to short periods of resistance training in young untrained men (Kraemer *et al.*, 1998, Kraemer *et al.*, 2005, Hansen *et al.*, 2001) or even long term trained weight lifters (Hakkinen *et al.*, 1988, Ahtianen *et al.*, 2003) at various ages (Kraemer *et al.*, 1999, Mc Call *et al.*, 1999, Hakkinen *et al.*, 2000). The response of GH to exercise in children seems to be related to the pubertal and

maturity stage of the individual (Bouix *et al.*, 1994, Marin *et al.*, 1994, Wirth and Gieck, 1994). Most studies in children and adolescents report resting plasma GH levels (Cacciari *et al.*, 1990, Mero *et al.*, 1990, Zakas *et al.*, 1994, Tsolakis *et al.*, 2003a,b, 2006), while there is a scarcity of data for the GH responses to exercise (Adiyaman *et al.*, 2004; Eliakim *et al.*, 2006).

### **Influence of resistance training on T, GH and neuromuscular performance in pre-pubescent, pubescent and adolescent individuals**

Since resistance training appears to increase strength in all ages, there appears to be some rationale for recommending resistance training as a means of improving sports performance even during preadolescence. There is insufficient evidence to state unequivocally that resistance training will have a positive effect on sports performance during preadolescence (Blimkie, 1993). Improvements in sport performance during preadolescence after different forms of resistance training have also been inferred from more indirect evidence, e.g. improvement in vertical jumping performance (Mero *et al.*, 1989, 1990a, b, Tsolakis *et al.*, 2003a, b, 2006 *in press*). However, even though there is a positive correlation between motor fitness and sports performance the results do not provide unequivocal evidence of a positive effect of resistance training on sports performance during preadolescence years.

The influence of various forms of sports training on the endocrine system has been studied mainly in adults involved in heavy training (Kraemer and Rogol, 2005). In fact there are very few data on the relationship between exercise and hormonal levels related to growth and maturation in sedentary pre-pubertal, pubertal and post-

pubertal children (Fahey et al., 1976, Zakas et al., 1993, Tsolakis et al., 2000, 2004). On the other hand more data exist for athletes in the pre-pubertal (Mero et al., 1989, 1990, a, b, Cacciari et al., 1990, Dally et al., 1998, Tsolakis et al., 2006) and post-pubertal period (Fry et al., 1993 Steinacker et al., 1993, Kraemer et al., 1992c, Gorostiaga et al., 1999, 2003).

Specifically T and FAI mean values have been reported to increase (Mero et al., 1990, Zakas et al., 1994, Tsolakis et al., 2000, 2004, 2006), or remain stable Dally et al., (1998) following two months to one year training. Some of these studies reported conflicting data on GH levels (Mero et al., 1990a, b, Zakas et al., 1994, Tsolakis et al., 2006). However most of these studies have suffered from methodological problems such as the use of small samples sizes that limit their external validity. The impact of acute exercise on the hypopituitary axis of youth has not been widely studied and there is not enough data to draw safe conclusions (Kraemer et al., 1992b).

#### **Effects of short and long term resistance training**

The effects of short term (2 months) resistance training on hormonal concentrations has been examined in two groups of untrained pre-pubertal and pubertal boys (Tsolakis et al., 2000). The mean post - training T concentration of the pre-pubertal group increased by 123%, whereas the respective increase in the pubertal group was 32% (Tsolakis et al., 2000). Another study published recently (Tsolakis et al., 2004), investigated the influence of two months of resistance training followed by two months of detraining on strength adaptations and selected hormones (T, SHBH) in sedentary pre-pubertal boys. Significant post training isometric strength gains (17.5%) and

increases in T and FAI ( $p < 0.05-0.001$ ) were observed in the experimental group. Detraining resulted in a significant loss (9.5%,  $p < 0.001$ ) of isometric strength whereas the hormonal parameters remained unchanged. The resistance training-induced strength changes were not correlated with changes in the anabolic and androgenic activity. Similar results were observed after six weeks of heavy resistance training in the dynamic strength of upper extremity (23%) and leg extensors (12%) in male 14-16 year old handball players. Moreover a significant increase was observed in throwing velocity in the resistance training group. Muscle hypertrophy did not accompany the strength gains, suggesting specific neurological adaptations (Gorostiaga et al., 1999).

The addition of either high or mild intensity cycloergometer training to three month sessions whole body calisthenics significantly increased T and GH levels in pubertal and post-pubertal sedentary boys but not in the pre-pubertal group (Zakas et al., 1994). An intensive gymnastic training program (3 h per day) performed by prepubertal male gymnasts was not adequate to alter serum T concentrations over a 10 month period. A possible explanation for this result is that the gymnasts were well adapted to the training (Daly et al., 1998). On the other hand, eleven weeks of soccer training combined with a program of explosive strength training resulted in significant increases in vertical jumping performance (14%) and in resting total T concentrations (7.5%) without any significant correlations between changes in performance and T (Gorostiaga et al., 2003).

The mean serum testosterone concentration increased significantly after 1 year of specific training in four groups of pre-pubescent athletes (sprinters, tennis

players, weight lifters and endurance runners), while GH remained unaltered. Additionally the experimental group increased speed and speed strength. Significant correlations were observed between the relative changes in testosterone, GH and the relative changes in speed. The increased anabolic activity had positive effects on physical performance and trainability at the early stages of puberty (*Mero et al., 1990*).

#### **Influence of specific sports training on hormonal levels**

There is a scarcity of data regarding the effects of specific sports training on hormonal levels in young athletes. *Mero et al., (1989)*, compared the neuromuscular, metabolic and hormonal profiles of trained pre-pubescent tennis players and an untrained group. Drop jump height was significantly greater ( $p < 0.05$ ) in the tennis players than in the control boys. However, there were non significant differences between the group serum hormone levels (*Mero et al., 1990*). Plasma levels of T, SHBG and GH were not different between of junior weight lifters, endurance and sprint runners. Interestingly, T concentration was significantly correlated with jumping performance. It was suggested that the training background and the advanced biological maturation of the athletes affected their strength capacity (*Mero et al., 1990*).

Significant differences were observed in plasma levels of dehydroepiandrosterone sulphate (DHEA-S) and T between pre-pubescent and pubescent football players and untrained controls (*Cacciari et al., 1990*). Plasma levels of T were significantly lower in the pre-pubescent group of athletes and significantly higher in the pubescent group compared to those of the controls. The higher values in pubertal subjects were justified by their advanced

maturation level compared to controls, while in pubertal subjects training may produce a decrease in plasma T as in adults (*Lauro et al., 1984, Hackney et al., 1988*).

*Tsolakis et al., (2003b)* demonstrated the effects of the specificity of training stimulus on the hormonal adaptations among pre-pubertal athletes of different sports. Figures 1 and 2 show the basal concentrations of T, GH, SHBG and FAI in 11-13 years old athletes of various sports (handball, rowing, running, basketball, swimming, weight-lifting and fencing), compared with a control group of untrained children. As can be seen, there are differences in hormonal concentrations between sports, probably because of differences in the content of training programs. The low T and GH in the weight lifting group was not expected, but may possibly be explained by their lower level of maturation compared to the other groups.

#### **Hormonal indices of overtraining and overreaching in young athletes**

As also observed in adult studies, the impact of very intense short duration in adolescent athletes of various sports result in significant hormonal changes, indicating a state of overreaching (*Fry et al., 1993, Steinacker et al., 1993, Gorostiaga et al., 1999*). Seven days of high volume resistance training sessions resulted in significant decreases in post exercise T levels in 28 junior elite weight lifters, indicating an impending overtraining syndrome (*Fry et al 1993*) since the high load of training may elicit an excessive endocrine demand in post-pubertal handball players (*Gorostiaga et al., 1999*). Significant increases in T and GH were reported after an acute weight lifting program in the most experienced athletes of the same study, suggesting that training experience in elite adolescent weightlifters is most influential on the hypopituitary-

gonadal training adaptations (Kraemer et al., 1992c). T and free T significantly decreased by 20 and 22% ( $p < 0.05$ ) after 16 days of high volume intensive rowing training in adolescent rowers. When the training volume decreased by 24% and the intensity was more than double T and free T increased significantly by 23 and 34.5% ( $p < 0.05$ ) respectively (Steinacker et al., 1993).

### Conclusions

The bulk of the available evidence would suggest that resistance training in children may result in an increase in hormones influencing anabolic activity and has a positive effect in physical performance and trainability even at the early stages of puberty. However, these changes are not always evident, probably due to the different characteristics of the resistance training programs across studies. It seems that there is an “intensity and volume threshold” of resistance training in order to have changes in hormonal concentrations that override the rapid growth process of puberty. Further longitudinal studies are needed to examine the influence of resistance training programs on the mechanisms that control growth and maturation.

### References

- Adiyaman, P., Ocal, G., Berberoglu, M., Evliyaoglu, O., Aycan, Z., Cetinkaya, E., Bulca, Y., Ersoz, G. and Akar, N. 2004. Alterations in serum growth hormone (GH)/GH dependent ternary complex components (IGF-I, IGFBP-3, ALS, IGF-I/IGFBP-3 molar ratio) and the influence of these alterations on growth pattern in female rhythmic gymnasts. *Journal of Pediatric Endocrinology and Metabolism*, **17**(6): 895-903.
- Adlercreutz, H., Harkonen, M., Kuoppasalmi, K., Tikkanen, H., Remes, K., Dessypris, A., and Karvonen, J. 1986. Effect of training on plasma anabolic and catabolic steroid hormones and their response during physical exercise. *International Journal of Sports Medicine*, **7**: S27-S28.
- Ahtiainen, J.P., Pakarinen, A., Kraemer, W.J., and Hakkinen, K. 2003. Muscle hypertrophy, hormonal adaptations and strength development during strength training in strength-trained and untrained men. *European Journal of Applied Physiology*, **89**: 555-563.
- Anderson, D.C. 1974. Sex hormone binding globulin. *Clinical Endocrinology*, **3**: 69-96.
- Araki, I., Harada, Y., and Kuno, M. 1991. Target-dependent hormonal control of neuron size in the rat spinal nucleus of the bulbocavernosus. *Journal of Neurosciences*, **11**: 3025-3033.
- BASES. 2004. Position statement on guidelines for resistance exercise in young people. *Journal of Sports Sciences*, **22**: 383-390.
- Blimkie, C.J.R., Ramsay, J., Sale, D., and Mac Dougall, D., and Smith, K. 1989. Effects of 10 weeks of resistance training on strength development on pre-pubertal boys. In: S. Oseid and K-H Carlsen (Eds), *Children and Exercise XIII*, 183-197. Champaign, IL: Human Kinetics.
- Blimkie, C.J.R. (1993). Resistance training during preadolescence. *Sports Medicine*, **15**(6): 389-407.
- Blimkie, C.J.R. and Bar-Or, O. 1996. Trainability of Muscle Strength, Power and Endurance during Childhood. In: *The child and adolescent athlete*. O Bar-Or (ed) Blackwell Science, 113-129.
- Boisseau, N., and Delamarche, P. 2000. Metabolic and Hormonal Responses to Exercise in children and Adolescents. *Sports Medicine*, **6**: 405-422.
- Bouix, O., Brun, J.F., Fedou, C., Raynard, E., Lenoir, V., and Orsetti, A. 1994. Plasma beta-endorphin corticotrophin and growth hormone to exercise in pubertal and pre-pubertal children. *Hormone and Metabolic Research*, **26**(4): 195-199.
- Cacciari, E.L., Mazzanti, L., Tassinari, R., Bergamaschi, C., Magnani, F., Zappulla, G., Nanni, C., Cobianchi, T., Ghini, R., and Tani, G. 1990. Effects of sport (football) on growth: Auxological, anthropometric and hormonal aspects. *European Journal of Applied Physiology*, **61**: 149-158.
- Crist, D.M., Peake, G.T., Lofffield, R.B., Kramer, J.C., and Egan, P.A. 1991. Supplemental growth hormones alter body composition, muscle protein metabolism and serum lipids in fit adults: characterization of dose-dependent and response-recovery effects. *Mechanisms of Ageing and Development*, **58**: 191-205.
- Dally, R.M., Rich, P.A., and Klein, R. 1988. Hormonal responses to physical training in high-level peripubertal male gymnasts. *European Journal of Applied Physiology*, **79**: 74-81.
- Eliakim, A., Nemet, D., Zaldivar, F., McMurray, R.G., Culler, F.L., Galassetti, P. and Cooper, D.M. 2006. Reduced exercise-associated response of the GH-IGF-I axis and catecholamines in obese children and adolescents. *Journal of Applied Physiology*, **100**(5): 1630-7.
- Fahey, T.D, Del Valle-Zuris, A., Ochisen, G., Tried, M., and Seymour, J. 1979. Pubertal stage differences in hormonal and hematological responses to maximal exercise in males. *Journal of Applied Physiology*. **46** (4): 823-827.
- Faigenbaum, A.D., Zaichkowsky, L.D., Wescott, W.L., Micheli, L., and Fehlandt, A.F. 1993. The effects



- of twice a week strength training program on children. *Paediatric Exercise Science*, **5**: 339-346.
- Falk, B. and Tenenbaum, G. 1996. The effectiveness of resistance training in children: a meta analysis. *Sports Medicine*, **22**: 176-186.
- Fleck, S.J. and Kraemer, W.J. 1987. *Designing resistance training programs*. Champaign, IL: Human Kinetics.
- Florini, J.R., Ewton, D.Z., and Coolican, S.A. 1996. Growth hormone and the insulin-like growth factor system in myogenesis. *Endocrine Reviews*, **17**(5): 481-517.
- Fry, A.C., Kraemer, W.J., Stone, M.H., Warren, B.J., Kearney, J.T., Fleck, S.J., and Weseman, C.A. 1993. Endocrine and performance responses to high volume training and amino acid supplementation in elite junior weight lifters. *International Journal of Sports Nutrition*, **3**: 306-322.
- Fry, A.C., Kraemer, W.J., Stone, M.H., Koziris, L.P., Thrush, J.T., and Fleck, S.J. 2000. Relationships between serum testosterone, cortisol and weightlifting performance. *Journal of Strength and Conditioning Research*, **14**(3): 338-343.
- Fukunaga, T., Funato, K., and Ikegawa, S. 1992. The effects of resistance training on muscle area and strength in pre-pubescent age. *The Annals of Physiological Anthropology*, **11**: 357-364.
- Gillam, G.M. 1981. Effects of frequency of weight training on muscle strength enhancement. *Journal of Sports Medicine*, **21**: 432-436.
- Gorostiaga, E.M., Izquierdo, M., Iturralde, P., Ruesta, M., and Ibanez, J. 1999. Effects of heavy resistance training on maximal and explosive force production, endurance and serum hormones in adolescent handball players. *European Journal of applied Physiology*, **80**: 485-490.
- Gorostiaga, E.M., Izquierdo, M., Ruesta, M., Iribarren, J., Gonzalez-Badillo, J.J., and Ibanez, J. 2004. Strength training effects on physical performance and serum hormones in young soccer players. *European Journal of Applied Physiology*, **91**: 698-707.
- Hacney, A.C., Sinning, W.E., and Bruot, B.C. 1988. Reproductive hormonal profiles of endurance-trained and untrained males. *Medicine and Science in sports and Exercise*, **20**: 60-65.
- Hakkinen, K., Pakarinen, A., Allen, M., Kauhanen, H., and Komi, P.V. 1987. Relationships between training volume, physical performance capacity and serum hormone concentrations during prolonged training in elite weight lifters. *International Journal of Sports Medicine*, **8**(Suppl): 61-65.
- Hakkinen, K. 1989. Neuromuscular and hormonal adaptations during strength and power training. *Journal of Sports Medicine and Physical Phitness*, **29**: 9-24.
- Hakkinen, K., Mero, A., and Kauhanen, H. 1989. Specificity of endurance, sprint and strenght training on physical performance capacity in young athletes. *Journal of sports Medicine*, **29**: 27-35.
- Hakkinen, K., and Pakarinen, A. 1995. Acute hormonal responses to heavy resistance exercise in men and women at different ages. *International Journal of Sports Medicine*, **16**(8): 507-513.
- Hakkinen, K., Pakarinen, A., Allen, M., Kauhanen, H., and Komi, P.V. 1998. Neuromuscular and hormonal adaptations in athletes to strength training in 2 years. *Journal of Applied Physiology*, **65**: 2406-2412.
- Hakkinen, K., Pakarinen, A., Kraemer, W.J., Newton, R.U., and Allen, M. 2000. Basal concentrations and acute response of serum hormones and strength development during heavy resistance training in middle-aged and elderly men and women. *Journals of Gerontology. Series A, Biological Sciences and Medical Sciences*, **55**: B95-B105.
- Hansen, S., Kvorning, T., Kjaer, M., Sjogaard, G. 2001. The effect of short-term strength training on human skeletal muscle: the importance of physiologically elevated hormone levels. *Scandinavian Journal of sports Medicine*, **11**: 347-354.
- Hassan , S.E.A. 1991. Die trainierbarkeit der maxinmalkraft bei 7-bis 13 jahrigen kindern. *Leistungssport*, **5**: 17-24.
- Jorgensen, J.O.L., Vahl, N., Hansen, T.B., Fisher, S., Hagen, C., and Christiansen, J.S. 1996. Influence of growth hormone and androgens on body composition in adults. *Hormone Research*, **45**: 94-98.
- Kraemer, W.J. 1988. Endocrine responses to resistance exercise. *Medicine Science in sports and Exercise*, **20** (suppl), S152-157.
- Kraemer, W.J., Fry, A.C., Frykman, P.N., Conroy, B., and Hoffman, J. 1989 Resistance training and Youth. *Paediatric Exercise Science*. **1**: 336-350.
- Kraemer, W.J., Marchitelli, L., Gordon, S.E., Harman, E., Dziados, J.E., Mello, R., Frykman, P., McCurry, D., and Fleck, S.J. 1990. Hormonal and growth factor responses to heavy resistance exercise protocols. *Journal of Applied Physiology*, **69**: 1442-1450.
- Kraemer, W.J., Gordon, S.E., Fleck, S.J., Marchitelli, L.J., Mello, R., Dziados, J.E., Friendl, K., and Harman, E. 1991. Endogenous anabolic responses to heavy resistance exercise in males and females. *International Journal of Sports Medicine*. **12**(2): 228-235.
- Kraemer, W.J. 1992a. Hormonal Mechanisms Related to the Expression of Muscular Strength and Power. In: *Strength and Power in Sport*. P.V. Komi, (ed). Blackwell Scientific Publications, 64-76.
- Kraemer, W.J. 1992b. Endocrine responses and Adaptations to Strength Training In: *Strength and Power in Sport*. P.V. Komi, (ed). Blackwell Scientific Publications, 291-304.
- Kraemer, W.J., Fry, A.C., Warren, B.J., Stone, M.H., Fleck, S.J., J.T. Kearney, J.T., Conroy, B.P., Maresh, C.M., Weseman, C.A., Triplett, N.T., and Gordon, S.E., 1992c. Acute hormonal responses

- in elite junior weightlifters. *International Journal of Sports Medicine*, **13** (2): 103-109.
- Kraemer, W.J. and Fleck, S.J. 1993. *Strength training for young athletes*. Champaign, IL. Human Kinetics
- Kraemer, W.J., Staron, P.S., Hagerman, F.C., Hikida, R.S., Fry, A.C., Gordon, S.E., Nindl, B.C., Gotshalk, L.A., Volek, J.S., Marx, J.O., Newton, R.U., and Hakkinen, K. 1998. The effects of short-term resistance training on endocrine function in men and women. *European Journal of Applied Physiology*, **78**: 69-76.
- Kraemer, W.J., Hakkinen, K., Newton, R.U., Nindl, B.C., Volek, J.S., McCormick, M., Gotshalk, L.A., Gordon, S.E., Fleck, S.J., Campbell, W.W., Putukian, M., and Evans, W.J. 1999. Effects of heavy resistance training on hormonal response patterns in younger vs older men. *Journal of Applied Physiology*, **87**: 982-992.
- Kraemer, W.J. and Rogol, A.D. (2005). *The Endocrine System in Sports and Exercise*. Blackwell Scientific Publications.
- Kraemer, W.J., Nindl, B.C., and Gordon, S.E. (2005) Resistance Exercise: Acute and Chronic Changes in Growth Hormone Concentrations. In: *The Endocrine System in Sports and Exercise*. Kraemer, W.J and A.D., Rogol (eds). Blackwell Scientific Publications, 110-121.
- Kuoppasalmi, K., Naveri, H., Harkonen, M., and Adlerkreutz, H. 1980. Plasma cortisol, androstendione, testosterone, and luteinizing hormone in running exercise of different intensities. *Scandinavian Journal of Clinical and Laboratory Investigations*, **40**: 403-409.
- Kurtz, E.M., Sengelau, D.R., and Arnold, A.P. 1986. Androgens regulate the dendritic length of mammalian motoneurons in adulthood. *Science*, **232**: 395-398.
- Lauro, P., N. Bergamo, N., R. Bertini, R., S. Dazzi, S., A. Dorigni, A., G.B. Lanzingher, G.B., Merz, O., Sarti, F., Zanon, B., Passamani, P., Schlechter, P., and Furlanello, F. (1984). Variazioni di alcuni parametrici ormonali in atleti della 'Marciaonga'. *Med Sport*, **37** : 409-417.
- Lee, P.A. 1995. Physiology of puberty. In Becker KL, ed. *Principles and practice of endocrinology and metabolism*, 2<sup>nd</sup> ed. Philadelphia: Lipincott: 822-830.
- Mantzoros, C.S., Flier, J.S., and Rogol, A.D. (1997). A longitudinal assessment of hormonal and physical alterations during normal puberty in boys. V. Rising leptin levels may signal the onset of puberty. *Journal of Clinical Endocrinology and Metabolism*, **82**: 1066-1070.
- Marin, G., Domene, H.M., Barnes, K.M., Blackwell, B.J., Cassorla, F.G., and Cutler, G.B. 1994. The effects of estrogen priming and puberty on the growth hormone response to standardized treadmill exercise and arginine-insulin in normal girls and boys. *Journal of Clinical Endocrinology and Metabolism*, **79**(2): 537-541.
- Matsumoto, A. 1992. Sex steroid induction of synaptic reorganisation in the adult *Neuroendocrine brain Reviews* **3**: 287-306.
- McCall, G.E. Byrnes, W.C., Fleck, S.J. and Kraemer, W.J. 1999. Acute and chronic responses to resistance training designed to promote muscle hypertrophy. *Canadian Journal of Applied Physiology*. **24**: 96-107.
- Mero, A., Jaakola, L. and Komi, P.V. 1989. Neuromuscular, metabolic and hormonal profiles of young tennis players and untrained boys. *Journal of Sports Sciences*, **7**: 95-100.
- Mero, A., Kauhanen, H., Peltola, E., Vuorimaa, T. and Komi, P.V. 1990a. Physiological performance capacity in different pre-pubescent athletic groups. *Journal of Sports Medicine and Physical Fitness*, **30**: 57-66
- Mero A., Jaakola, L. and Komi, P.V. 1990b. Serum hormones and physical performance capacity in young boys athletes during a 1-year training period. *European Journal of Applied Physiology and Occupational Physiology*, **60**: 32-37.
- Mersch, F. and Stoboy, H. 1989. Strength training and muscle hypertrophy in children. In S. Oseid-K-H. Karlsen, (eds) *Children and Exercise XIII*, 165-183. Human Kinetics, Champaign IL.
- Moritani, T. 1992. Time course of adaptations during strength and power training. In: *Strength and Power in Sport*. P.V. Komi, ed. Blackwell Scientific Publications, 266-278.
- Moritani, T. 1993. Neuromuscular adaptations during the acquisition of muscle strength, power and motor tasks. *Journal of Biomechanics*, **26**: 95-107.
- Nielsen, B., Nielsen, K., Behrendt-Hansen, M. and Asmusen, E. (1980). Training of functional muscular strength in girls 7-19 years old. In K.Berg and B.D. Eriksson (eds) *Children and Exercise IX*, 69-78. Human Kinetics, Champaign, IL.
- Nindl, B.C., Kraemer, W.J., Marx, J.O., Tuckow, A.P. and Hymer, W.C. 2003. Growth hormone molecular heterogeneity and exercise. *Exercise and Sports Science Reviews*, **31**(4): 161-166.
- Ozmun, J.C., Mikesky, A.E., and Surburg, P.R. 1994. Neuromuscular adaptations following pre-pubescent strength training. *Medicine Science in Sports and Exercise*, **26**: 510-514.
- Pfeifer, R.D. and Francis, R.S. 1986. Effects of strength training on muscle development in pre-pubescent, pubescent and post pubescent males. *Physician and Sports Medicine*, **14**: 134-143.
- Ramsay, J.A., Blimkie, C.J.R., Smith, K., Garner, S., Mac Dougal S.D. and Sale, D.G. 1990). Strength training effects in pre-pubescent boys. *Medicine and Science in Sports and Exercise*, **22**: 605-614.
- Remes, K., Kuoppasalmi, K. and Adlerkreutz, H.. 1979. Effect of long term physical training on plasma testosterone, androstendione, luteinizing hormone and sex hormone binding globulin capacity. *Scandinavian Journal of Clinical Investigation*, **39**: 743-749.

- Roemmich, J.N. (2005). Growth, Maturation and Hormonal Changes During Puberty: Influence of Sport Training. In: *The Endocrine System in Sports and Exercise*. Kraemer, W.J and A.D., Rogol (eds). Blackwell Scientific Publications, 512-524.
- Roemmich, J.N. and Rogol, A.D. 1997. Exercise and growth hormone: does one affects other? *Journal of Pediatrics*, **131(1pt2)**: S75-S80.
- Rogol, A.D. 1994. Growth at puberty: interaction of androgen and growth hormone. *Medicine and Science in Sports and Exercise*, **26**: 767 –770.
- Rosner, W. 1991. Plasma steroid-binding proteins. *Endocrinology and Metabolism Clinics of North America*, **20(4)**: 697-720.
- Sailors, M. and Berg, K. 1987. Comparison of responses to weight training in pubescent boys and men. *Journal of Sports Medicine and Physical Fitness*, **27**: 30-37.
- Sale, D.G. 1989. Strength training in children. In C.V. Gisolfi and D.R. Lamb (eds) *Perspectives in Exercise Science and Sports Medicine*, Vol. 2 Youth and Exercise and Sport, 165-222. Benchmark Press, Indianapolis..
- Schmid, P., Push, H.H., Wolf, W., Pilger, E. and Pessenhofer, H. 1982. Serum FSH, LH, and testosterone in humans after physical exercise. *International Journal of Sports Medicine*, **3**: 84-89.
- Schurmeyer, T., Wickings, E.J., Freisheim, C.W. and Nieschlag, E. (1983) Saliva and serum testosterone following oral testosterone undecanoate administration in normal and hypogonadal men. *Acta Endocrinologica*, **102**: 456-462.
- Servedio, F.J., Bartels, R.L., Hamlin, R.L., Teske, D., Shafer, T. and Servedio, A. 1985. The effects of weight training using Olympic style lifts on various physiological variables in pre-pubescent boys. *Medicine and Science in sports and Exercise*, **17**: 288.
- Shephard, R.J. 1984. Physical activity and child health. *Sports Medicine*, **1**: 205-233.
- Siegel, J.A., Camaione, D.N. and Manfredi, T.G. (1989) The effects of upper body resistance training on pre-pubescent children. *Paediatric Exercise Science*, **1**: 145-154.
- Staron, R.S., Karapondo, D.L., Kraemer, W.J., Fry, A.C., Gordon, S.E., Falkel, J.E., Hagerman, F.C. and Hikida, R.S. 1994. Skeletal muscle adaptation during the early phase of heavy resistance training in men and women. *Journal of Applied Physiology*, **76**: 1247-1255.
- Steinacker, J.M., Laske, R., Hetzel, W.D., Lormes, W., Liu, Y. and Stauch, M. 1993. Metabolic and Hormonal Reactions During Training in Junior Oarsmen. *International Journal of Sports Medicine*, **14(Suppl)**, S24-S28.
- Tsolakis, C., Messinis, D., Stergioulas, A. and Dessypris A. 2000. Hormonal responses after strength training and detraining in pre-pubertal and pubertal boys. *Journal of Strength and Conditioning Research*, **14(4)**: 399-404.
- Tsolakis, C., Vagenas, G. and Dessypris, A. 2003a. Leptin, Testosterone, GH, body composition and neuromuscular performance in pre-pubescent athletes and untrained boys. *Journal of Strength and Conditioning Research*, **17**: 40-46.
- Tsolakis C, Xekouki, P., Kaloupsis, S., Karras, D., Messinis, D., Vagenas, G. and Dessypris, A. 2003b. The influence of exercise on growth hormone and testosterone in pre-pubertal and early pubertal boys. *Hormones*, **2(2)**: 13-112.
- Tsolakis, C., Vagenas, G. and Dessypris, A. 2004. Strength adaptations and hormonal responses to resistance training and detraining in preadolescent males. *Journal of Strength and Conditioning Research*. **18(3)**: 625-629.
- Tsolakis, C., Bogdanis, G.C., Vagenas, G. and Dessypris, A. 2006. Influence of a 12-month conditioning program on physical growth, serum hormones and neuromuscular performance of peripubertal male fencers. *Journal of Strength and Conditioning Research* (in press).
- Viru, A. and Viru, M. 2005. Resistance Exercise and Testosterone. In: *The Endocrine System in Sports and Exercise*. Kraemer, W.J and A.D., Rogol (eds). Blackwell Scientific Publications, 319-338.
- Viru, A.A. 1985. *Hormones in Muscular Activity*. I. Hormonal Ensemble in Exercise. CRC Press, Boca Raton, FL: 87.
- Weiss, R.E. and Refetoff, S. 1996. Effect of thyroid hormone on growth: lessons from the syndrome of resistance to thyroid hormone. *Endocrinology and Metabolism Clinics of North America*, **25(3)**: 719-730.
- Weltman, A., Janney, C., Rians, C.B., Strand, K., Berg, B., Tippitt, S., Wise, J., Cahill, B. and Katch, F. 1986. The effects of hydraulic resistance strength training in pre-pubertal males. *Medicine and Science in Sports and Exercise*, **18**: 629-638.
- Wirth, V.J. and Gieck, J. 1994. Growth hormone: Myths and misconceptions. *Journal of sports Rehabilitation*, **5(3)**: 244-250.
- Zakas, A., Mandroukas, K., Karamouzis, M. and Panagiotopoulou, G. 1994. Physical training, growth hormone, and testosterone levels and blood pressure in pre-pubertal, pubertal and adolescent boys. *Scandinavian Journal of Medicine and Science in Sports*. **4**: 113-118.